

Lemma: Let  $G$  be a finite group,  $P$  a Sylow  $p$ -subgroup,  $H \subset G$  a  $p$ -group. If  $H \subset N_G(P)$  then

$H \subset P$ , and if  $H$  is a Sylow  $p$ -subgroup then  $H = P$ .

Notation: For  $G$  group, let  $S = \mathcal{P}(G)$ , consider the action  $* : G \times S \rightarrow S$ . We set:

$$(x, A) \mapsto xAx^{-1}$$

$C(A) := G * A = \{xAx^{-1} \mid x \in G\}$ , now for  $H \subset G$  we can restrict this action to:

$*|_{G \times C(H)} : G \times C(H) \rightarrow C(H)$ , making  $C(H)$  into a  $G$ -set.

Lemma: Let  $G$  be a finite group,  $P$  a Sylow  $p$ -subgroup,  $H \subset G$  a  $p$ -group. Then:

1.  $C(P)$  consists of Sylow  $p$ -subgroups.

2. The conjugacy class  $C(P)$  is an  $H$ -set by conjugation.

3. If  $T$  is a fixed point under the  $H$ -action, namely:

$$T \in F_H(C(P)) = \{w \in C(P) \mid xwx^{-1} = w \text{ for all } x \in H\}, \text{ then } H \leq T.$$

4. If  $H$  is a Sylow  $p$ -subgroup then under the  $H$ -action,  $H$  is the only possible fixed point:

$$F_H(C(P)) \leq \{H\}.$$

Theorem: (Second Sylow Theorem) Let  $G$  a finite group,  $p \in \mathbb{N}^+$  a prime dividing  $|G|$ . Then all

Sylow  $p$ -subgroups are conjugate:  $P \in \text{Syl}_p(G)$  then  $C(P) = \text{Syl}_p(G)$ .

Theorem: (Third Sylow Theorem) Let  $G$  a finite group,  $p \in \mathbb{N}^+$  a prime dividing  $|G|$ . Let  $P$  be

a Sylow  $p$ -subgroup, then:

$$(i) [G : N_G(P)] = |\text{Syl}_p(G)|.$$

$$(ii) |\text{Syl}_p(G)| \text{ divides } |G|$$

$$(iii) |\text{Syl}_p(G)| \equiv 1 \pmod{p}$$

$$(iv) |\text{Syl}_p(G)| \text{ divides } [G : P], \text{ namely if } |G| = p^m n \text{ with } p \nmid n \text{ then } |\text{Syl}_p(G)| \mid n.$$

Theorem: (Fourth Sylow Theorem) Let  $G$  a finite group,  $p \in \mathbb{N}^+$  a prime dividing  $|G|$ . Let  $H$  be a  $p$ -subgroup of  $G$ . Then  $H$  lies in some Sylow  $p$ -subgroup of  $G$ .

Proof: (Second, Third, and Fourth Sylow Theorems) Let  $H$  be a subgroup of  $G$  that is a  $p$ -group,

and  $P$  is a Sylow  $p$ -subgroup of  $G$ . By the Lemma above,  $C(P) \subseteq \text{Syl}_p(G)$  is an  $H$ -set

under conjugation. By the Orbit Decomposition Theorem:

$$|C(P)| = |F_H(C(P))| + \sum_{T \in G^*} [H : H_T].$$

If  $T \in G^*$  then  $H_T \not\subseteq H$  so  $p$  divides  $[H : H_T]$ , so  $|C(P)| \equiv |F_H(C(P))| \pmod{p}$ .

If  $H = P$ , by the Lemma above we have  $F_P(C(P)) \subseteq \{P\}$ . Since  $xPx^{-1} = P$  for all

$x \in P$  we have  $F_P(C(P)) = \{P\}$ . Now by the above:

$$|C(P)| \equiv |F_P(C(P))| \equiv 1 \pmod{p}.$$

If  $H$  is a Sylow  $p$ -subgroup of  $G$ , not necessarily  $P$ , then:

$|F_H(C(P))| \equiv |C(P)| \equiv 1 \pmod{p}$ . Hence  $F_H(C(P))$  is not empty, so by the Lemma above,  $F_H(C(P)) = \{H\}$ . In particular  $H \in F_H(C(P)) \subseteq C(P)$ . Since  $H$  is any Sylow  $p$ -subgroup, then  $Syl_p(G) \subseteq C(P)$ . So  $Syl_p(G) = C(P)$ . This proves the Second Sylow Theorem. Hence:  $|Syl_p(G)| \equiv |C(P)| \equiv 1 \pmod{p}$ , also  $|Syl_p(G)| = |C(P)| = [G : N_G(P)]$ ,

and  $|G| = [G : N_G(P)] \cdot |N_G(P)| = |Syl_p(G)| \cdot |N_G(P)|$  so  $|Syl_p(G)|$  divides  $|G|$ . Moreover

since  $P \subseteq N_G(P)$  we have:  $[G : P] = [G : N_G(P)] [N_G(P) : P] = |Syl_p(G)| \cdot [N_G(P) : P]$  so

$|Syl_p(G)|$  divides  $[G : P]$ , proving the Third Sylow Theorem.

If  $H$  is a  $p$ -subgroup of  $G$ , not necessarily a Sylow  $p$ -subgroup, we still have:

$$|F_H(C(P))| \equiv |C(P)| \equiv 1 \pmod{p}.$$

Thus there exists  $T \in F_H(C(P)) \subseteq C(P) = Syl_p(G)$ , so by the Lemma above  $H \subseteq T$ ,

and  $T$  is a Sylow  $p$ -subgroup. So  $H$  is contained in a Sylow  $p$ -subgroup, proving the

Fourth Sylow Theorem. □.