

**Math 115A**  
**Linear Algebra**

**Discussion 6**

**Problem 1(★).**

An *elementary matrix* is a matrix obtained from the identity by performing one elementary row operation.

- Denote by  $T_{ij}$  the elementary matrix obtained by exchanging the  $i$ -th and  $j$ -th rows. Write  $T_{ij}$  in matrix form. Compute  $\det(T_{ij})$ . Prove that  $\det(T_{ij}^t) = \det(T_{ij})$ . Prove that  $T_{ij}^{-1} = T_{ij}$ .
- Denote by  $D_i(m)$  the elementary matrix obtained by multiply the  $i$ -th row by a scalar  $m$ . Write  $D_i(m)$  in matrix form. Compute  $\det(D_i(m))$ . Prove that  $\det(D_i(m)^t) = \det(D_i(m))$ . Prove that  $D_i(m)^{-1} = D_i(1/m)$ .
- We denote by  $L_{ij}(m)$  the elementary matrix obtained by adding to the  $i$ -th row the  $j$ -th row multiplied by a scalar  $m$ . Write  $L_{ij}(m)$  in matrix form. Compute  $\det(L_{ij}(m))$ . Prove that  $\det(L_{ij}(m)^t) = \det(L_{ij}(m))$ . Prove that  $L_{ij}(m)^{-1} = L_{ij}(-m)$ .

**Problem 2.**

A matrix  $M \in M_{n \times n}(\mathbb{C})$  is called *nilpotent* if  $M^k = O$  for some positive integer  $k$ . Prove that if  $M$  is nilpotent then  $\det(M) = 0$ .

**Problem 3.**

A matrix  $M \in M_{n \times n}(\mathbb{C})$  is called *skew-symmetric* if  $M^t = -M$ . Prove that if  $M$  is skew-symmetric and  $n$  is odd then  $M$  is not invertible. What happens if  $n$  is even? Give examples if possible.

**Problem 4.**

A matrix  $Q \in M_{n \times n}(\mathbb{R})$  is called *orthogonal* if  $QQ^t = I_n$ . Prove that if  $Q$  is orthogonal then  $\det(Q) \in \{-1, 1\}$ .

**Problem 5(★).**

Let  $M \in M_{n \times n}(\mathbb{C})$ , define the matrix  $\overline{M}$  via  $(\overline{M})_{ij} = \overline{M_{ij}}$  for all  $i, j \in \{1, \dots, n\}$ .

- Prove that  $\det(\overline{M}) = \overline{\det(M)}$ .
- Prove that  $\overline{M^t} = (\overline{M})^t$ . Define  $M^* = \overline{M^t}$ .
- A matrix  $Q \in M_{n \times n}(\mathbb{C})$  is called *unitary* if  $QQ^* = I_n$ . Prove that if  $Q$  is unitary then the modulus of the complex number  $\det(Q)$  is 1, that is,  $|\det(Q)| = 1$ .

**Problem 6.**

A matrix  $A \in M_{n \times n}(\mathbb{F})$  is called *lower triangular* if  $A_{ij} = 0$  whenever  $1 \leq i < j \leq n$ . Let  $A$  be lower triangular, describe  $\det(A)$  in terms of the entries of  $A$ . Prove your claim.

**Problem 7.**

Let  $A, B \in M_{n \times n}(\mathbb{F})$ . Prove that if  $A$  is similar to  $B$  then  $\det(A) = \det(B)$ .