

Math 115A
Linear Algebra

Discussion for May 9-13, 2022

Problem 1.

Let V be a finite dimensional vector space over \mathbb{R} . Show that if $\dim(V)$ is odd, then every $T \in \mathcal{L}(V)$ has an eigenvalue.

Problem 2.

Let V be a finite dimensional vector space over \mathbb{R} and $T \in \mathcal{L}(V)$ has no real eigenvalues. Prove that every T -invariant subspace of V has even dimension.

Problem 3.

Let V be a finite dimensional vector space over \mathbb{R} . Prove that every $T \in \mathcal{L}(V)$ has an invariant subspace of dimension one or two.

Problem 4.

Let V be a vector space over \mathbb{F} of dimension n . Suppose that $T \in \mathcal{L}(V)$ has n distinct eigenvalues.

- (a) Prove that T has n distinct eigenvectors forming a basis of V .
- (b) Prove that if $S \in \mathcal{L}(V)$ has the same eigenvectors as T (but not necessarily the same eigenvalues) then $ST = TS$.

Problem 5.

Let V be a vector space over \mathbb{F} of dimension n . Suppose that $T \in \mathcal{L}(V)$ is such that all subspaces of V of dimension $n - 1$ are T -invariant. Show that T is a scalar multiple of the identity operator.

Problem 6(★).

Let V be a vector space over \mathbb{F} of dimension n , let $T \in \mathcal{L}(V)$, let β be an ordered basis of V . The *determinant* of T , denoted $\det(T)$, is defined as $\det(T) = \det([T]_\beta)$.

- (a) Prove that the determinant of T is independent of the choice of β . Namely, prove that if β and γ are two ordered bases of V , then $\det([T]_\beta) = \det([T]_\gamma)$.
- (b) Prove that T is invertible if and only if $\det(T) \neq 0$.
- (c) Prove that if T is invertible, then $\det(T^{-1}) = \det(T)^{-1}$.
- (d) Prove that if $S \in \mathcal{L}(V)$ then $\det(TS) = \det(T) \det(S)$.
- (e) Prove that if $\lambda \in \mathbb{F}$ then $\det(T - \lambda I_V) = \det([T]_\beta - \lambda I_n)$.