

**Math 33A**  
**Linear Algebra and Applications**

**Discussion for February 28-March 4, 2022**

**Problem 1.**

Consider the  $n \times n$  matrix

$$J_n(k) = \begin{bmatrix} k & 1 & 0 & \cdots & 0 & 0 \\ 0 & k & 1 & \cdots & 0 & 0 \\ 0 & 0 & k & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & k & 1 \\ 0 & 0 & 0 & \cdots & 0 & k \end{bmatrix}$$

(with all  $k$ 's on the diagonal and 1's directly above), where  $k$  is an arbitrary constant. Find the eigenvalue(s) of  $J_n(k)$ , and determine their algebraic and geometric multiplicities.

**Solution:** Since  $J_n(k)$  is upper triangular, its eigenvalues are its diagonal entries, so it has  $k$  as its single eigenvalue, with algebraic multiplicity  $n$ . Since

$$E_k = \ker(J_n(k) - kI_n) = \ker \left( \begin{bmatrix} k & 1 & 0 & \cdots & 0 & 0 \\ 0 & k & 1 & \cdots & 0 & 0 \\ 0 & 0 & k & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & k & 1 \\ 0 & 0 & 0 & \cdots & 0 & k \end{bmatrix} \right) = \text{span}(\vec{e}_1)$$

then  $\dim(E_k) = 1$ , so the geometric multiplicity of  $k$  is 1.

**Problem 2.**

Are the following matrices similar?

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

**Solution:** No, since  $A^2 = 0$  but  $B^2 \neq 0$ . If  $A$  was similar to  $B$ , then we would have an invertible matrix  $S$  satisfying  $B = S^{-1}AS$ , and thus  $0 \neq B^2 = S^{-1}A^2S = 0$  gives a contradiction.

**Problem 3.**

Consider a nonzero  $3 \times 3$  matrix  $A$  such that  $A^2 = 0$ .

- (a) Show that the image of  $A$  is a subspace of the kernel of  $A$ .
- (b) Find the dimensions of the image and kernel of  $A$ .
- (c) Pick a nonzero vector  $v_1$  in the image of  $A$ , and write  $\vec{v}_1 = A\vec{v}_2$  for some  $\vec{v}_2$  in  $\mathbb{R}^3$ . Let  $\vec{v}_3$  be a vector in the kernel of  $A$  that fails to be a scalar multiple of  $\vec{v}_1$ . Show that  $\mathfrak{B} = \{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$  is a basis of  $\mathbb{R}^3$ .
- (d) Find the matrix  $B$  of the linear transformation  $T(\vec{x}) = A\vec{x}$  with respect to basis  $\mathfrak{B}$ .

**Solution:**

- (a) Let  $\vec{v} \in \text{im}(A)$ , so that there is a vector  $\vec{w} \in \mathbb{R}^3$  with  $\vec{v} = A\vec{w}$ . Now  $A\vec{v} = A(A\vec{w}) = A^2\vec{w} = 0\vec{w} = \vec{0}$ , so  $\vec{v} \in \text{ker}(A)$ . Thus  $\text{im}(A) \subset \text{ker}(A)$ .
- (b) By the above,  $\dim(\text{im}(A)) \leq \dim(\text{ker}(A))$ . Since  $A$  is non zero, then we have at least one non zero vector in the image of  $A$ , so  $\dim(\text{im}(A)) \geq 1$ . By the rank nullity Theorem we have  $\dim(\text{im}(A)) + \dim(\text{ker}(A)) = 3$ . Thus since the dimensions are integers, the only possibility is  $\dim(\text{im}(A)) = 1$  and  $\dim(\text{ker}(A)) = 2$ .
- (c) We have three non zero vectors, so to prove that they are a basis of  $\mathbb{R}^3$  it is enough to prove that they are linearly independent. Suppose we have a relation  $c_1\vec{v}_1 + c_2\vec{v}_2 + c_3\vec{v}_3 = \vec{0}$  for some real scalars  $c_1, c_2, c_3$ . Applying  $A$  to both terms of the equality we obtain  $\vec{0} = c_2A\vec{v}_2 = c_2\vec{v}_1$  so  $c_2 = 0$ , using that  $c_2, c_3$  are in  $\text{ker}(A)$ . Thus we have  $c_2\vec{v}_2 + c_3\vec{v}_3 = \vec{0}$ . Since  $\vec{v}_2$  and  $\vec{v}_3$  are linearly independent by construction, we have  $c_2 = c_3 = 0$ . Hence  $\vec{v}_1, \vec{v}_2, \vec{v}_3$  are linearly independent.
- (d) We have

$$B = [[A(\vec{v}_1)]_{\mathfrak{B}} \quad [A(\vec{v}_2)]_{\mathfrak{B}} \quad [A(\vec{v}_3)]_{\mathfrak{B}}] = [[\vec{0}]_{\mathfrak{B}} \quad [\vec{v}_1]_{\mathfrak{B}} \quad [\vec{0}]_{\mathfrak{B}}] = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

**Problem 4.**

If  $A$  and  $B$  are two nonzero  $3 \times 3$  matrices such that  $A^2 = B^2 = 0$ , is  $A$  necessarily similar to  $B$ ?

**Solution:** Yes. Using the previous problem, we can find a basis  $\mathfrak{A}$  of  $\mathbb{R}^3$  such that  $A$  is similar to  $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ , and we can also find a basis  $\mathfrak{B}$  of  $\mathbb{R}^3$  such that  $B$  is similar to  $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ . Thus, since  $A$  and  $B$  are both similar to the same matrix, they are similar to each other.

**Problem 5.**

For the matrix

$$A = \begin{bmatrix} 1 & -2 & 1 \\ 2 & -4 & 2 \\ 3 & -6 & 3 \end{bmatrix},$$

find an invertible matrix  $S$  such that

$$S^{-1}AS = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

**Solution:** Note that  $A^2 = 0$ , so we can use the method given above. We know that the vector  $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$  is in the image of  $A$ , since it is the first column of  $A$ . Thus the way we obtain it is multiplying  $A$  by the first vector of the standard basis, namely  $\vec{v}_1 = A\vec{e}_1$ , so we set  $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ . For our last element of the basis, we need a vector in the kernel of  $A$  that is not a scalar multiple of  $\vec{v}_1$ . Since

$$\text{rref}(A) = \begin{bmatrix} 1 & -2 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

we have the two relations  $\vec{v}_2 = -2\vec{v}_1$  and  $\vec{v}_3 = \vec{v}_1$ , giving the vectors  $\begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$  and  $\begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$  in the kernel of  $A$ . Neither of them is a scalar multiple of  $\vec{v}_1$ , so we can set  $\vec{v}_3 = \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix}$ .

Now

$$\mathfrak{B} = \left\{ \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} \right\}$$

is a basis of  $\mathbb{R}^3$  such that the linear transformation associated to  $A$  in  $\mathfrak{B}$  is  $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ .

In particular

$$\begin{bmatrix} 1 & 1 & 2 \\ 2 & 0 & 1 \\ 3 & 0 & 0 \end{bmatrix}^{-1} \begin{bmatrix} 1 & -2 & 1 \\ 2 & -4 & 2 \\ 3 & -6 & 3 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 2 & 0 & 1 \\ 3 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

as desired.

**Problem 6.**

Consider an  $n \times n$  matrix  $A$  such that  $A^2 = 0$ , with  $\text{rank}(A) = r$  (above we have seen the case  $n = 3$  and  $r = 1$ ). Show that  $A$  is similar to the block matrix

$$B = \begin{bmatrix} J & 0 & \cdots & 0 & \cdots & 0 \\ 0 & J & \cdots & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & & \vdots \\ 0 & 0 & \cdots & J & \cdots & 0 \\ \vdots & \vdots & & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & \cdots & 0 \end{bmatrix}, \quad \text{where } J = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

Matrix  $B$  has  $r$  blocks of the form  $J$  along the diagonal, with all other entries being 0. To show this, proceed as in the case above: Pick a basis  $\vec{v}_1, \dots, \vec{v}_r$  of the image of  $A$ , write  $\vec{v}_i = A\vec{w}_i$  for  $i = 1, \dots, r$ , and expand  $\vec{v}_1, \dots, \vec{v}_r$  to a basis  $\vec{v}_1, \dots, \vec{v}_r, \vec{u}_1, \dots, \vec{u}_m$  of the kernel of  $A$ . Show that  $\vec{v}_1, \vec{w}_1, \vec{v}_2, \vec{w}_2, \dots, \vec{v}_r, \vec{w}_r, \vec{u}_1, \dots, \vec{u}_m$  is a basis of  $\mathbb{R}^n$ , and show that  $B$  is the matrix of  $T(\vec{x}) = A\vec{x}$  with respect to this basis.

**Solution:** To show that  $\vec{v}_1, \vec{w}_1, \vec{v}_2, \vec{w}_2, \dots, \vec{v}_r, \vec{w}_r, \vec{u}_1, \dots, \vec{u}_m$  is a basis of  $\mathbb{R}^n$  it is enough to prove that there are  $n$  of them and that they are linearly independent.

Since  $\vec{v}_1, \dots, \vec{v}_r$  form a basis of the image of  $A$  we have  $\dim(\text{im}(A)) = r$ . Since  $\vec{v}_1, \dots, \vec{v}_r, \vec{u}_1, \dots, \vec{u}_m$  is a basis of the kernel of  $A$  then  $\dim(\text{ker}(A)) = r + m$ . Thus by the rank nullity Theorem we have  $n = \dim(\text{im}(A)) + \dim(\text{ker}(A)) = r + r + m = 2r + m$  so indeed there are  $n$  vectors in the list  $\vec{v}_1, \vec{w}_1, \vec{v}_2, \vec{w}_2, \dots, \vec{v}_r, \vec{w}_r, \vec{u}_1, \dots, \vec{u}_m$ .

To see that they are linearly independent, suppose we have a linear combination  $a_1\vec{v}_1 + b_1\vec{w}_1 + \cdots + a_r\vec{v}_r + b_r\vec{w}_r + c_1\vec{u}_1 + \cdots + c_m\vec{u}_m = \vec{0}$ . Applying  $A$  to both sides of the equality we obtain  $b_1\vec{v}_1 + \cdots + b_r\vec{v}_r = \vec{0}$  so  $b_1 = \cdots = b_r = 0$  since  $\vec{v}_1, \dots, \vec{v}_r$  are linearly independent. We then have  $a_1\vec{v}_1 + \cdots + a_r\vec{v}_r + c_1\vec{u}_1 + \cdots + c_m\vec{u}_m = \vec{0}$ , so  $a_1 = \cdots = a_r = c_1 = \cdots = c_m = 0$  since  $\vec{v}_1, \dots, \vec{v}_r, \vec{u}_1, \dots, \vec{u}_m$  are linearly independent.

What remains is to show that the matrix  $B$  is similar to  $A$  with respect to this change of basis. Note that for each  $i = 1, \dots, r$  the pair  $\vec{v}_i, \vec{w}_i$  will contribute with a block

$$J = \begin{bmatrix} [A(\vec{v}_i)]_{\{\vec{v}_i, \vec{w}_i\}} & [A(\vec{w}_i)]_{\{\vec{v}_i, \vec{w}_i\}} \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} b_{i,i} & b_{i,i+1} \\ b_{i+1,i} & b_{i+1,i+1} \end{bmatrix}$$

to the matrix  $B$ , these blocks having their diagonal coincide with the diagonal of  $B$ . Moreover, since  $A\vec{u}_j = \vec{0}$  for all  $j = 1, \dots, m$ , all the other entries of the matrix  $B$  are zero.